

Study of cirrus clouds and implications in the variability of laser propagation light and variability of fratricide effect

Angel Otarola^a and Paul Hickson^b

a) TMT International Observatory

b) University of British Columbia

June 30, 2017

AO4ELT5 Conference
Puerto de La Cruz, Tenerife, Spain

Motivation

- “AO observations (mainly LGS) are difficult to make and history [at some observatories] suggests that when atmospheric conditions are worse than median, the productivity is dramatically lower... And cirrus clouds can dramatically hurt AO.
- New generation large aperture telescopes are planning to make extensive use of multiple lasers to achieve diffraction limited imaging capabilities.
 - In the case of TMT, and for early science/operations phase, the plan is to dedicate the telescope observing time approximately as:
 - 40%, Visible seeing limited imaging/spectroscopy
 - **50%, Near Infrared diffraction limited imaging/spectroscopy**
 - 10% Mid Infrared programs
- Besides, it is of interest to have a better idea of what are the clear fraction and cirrus cloud properties (altitude and thickness) for the potential sites to host the TMT project.
 - Learning the seasonal changes in cirrus cloud and atmospheric aerosol is also of interest to know.

Goals

◆ Learn about:

- ◇ All year around fraction of **CLEAR** and **USABLE** nights in a year at various sites where Laser Systems to support Laser Guide Star (LGS) AO are operated.
- ◇ The statistics of altitude and thickness of cirrus clouds.
- ◇ The statistics of optical depth for cirrus clouds and atmospheric aerosols in the upper atmosphere.
- ◇ Changes in the these parameters as a function of month through the year.
- ◇ Effects of cirrus in the intensity of LGS


Previous Studies Usable Nights

- **Dr. Andre Erasmus and Collaborators (USABLE nights, **annual fraction**)**
 - GOES-8 satellite data consist of infra-red window (10.7 μ m) – for CLOUDS
 - GOES-8 satellite data ((6.7 μ m) for water vapour
 - Images every 3 hours (0245UT, 0545UT, 0845UT, 1145UT, 1445UT, 1745UT, 2045UT and 2345UT) :
 - Cerro Tololo/Cerro Pachón (CHILE) **78.2%**
 - La Silla/Las Campanas (CHILE) **83.1%**
 - Armazones/Paranal (CHILE) **90.5%**
 - Mount Graham/Kitt Peak (USA)
 - Maunakea (USA) **71.4%**

- **IAC team (Garcia-Gill et al., 2010) –from weather downtime of various telescopes**
 - Canary Islands (SPAIN) (69.7% – 79.3%) **72 %**


Data for this study

- We make use of observations using the CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) Sensor on board of the CALIPSO Satellite.




Parameter	Magnitude
Laser	Nd:YAG, diode-pumped, Q-switched frequency doubled
Wavelength	532 nm, 1064 nm
Pulse Energy	110 mJ/channel
PRR	20.25 Hz
Receiver diameter	1.0 m
Polarization	532 nm Lidar
Footprint/FOV	100 m / 130 urad
Vertical Resolution	30 – 60 m
Horizontal Resolution	5 km (above 8000 m) 333 m (below 800 m)

Examples of CALIPSO (A-Train Orbit) Data period 2006-2016



Maunakea











Canary Islands

Assumptions








- Clear fraction is that when no clouds were detected by the 532 nm Lidar in the CALIPSO satellite at any level between the altitude of the site and above.
- Usable time is defined as the Clear Fraction plus the existence of **Thin Cirrus** clouds with optical depths of up to 0.3 nepers (~ 0.28 mag/airmass)
 - Typical optical depth due to molecular scattering ~ 0.05
 - \rightarrow Minimum atmospheric transmission $\sim \exp(- [0.3+0.05]) \sim \mathbf{0.7}$
- Only night-time data used for the statistics (information from CALIOP for day time is available but not fully analyzed)

ANNUAL Trends in Clear (Blue) and Usable Time (red) for various sites of astronomical interest



- ORM Site: There is evidence the increase in usable nights → + Phase of NAO
- Chile: a clear trend in the usable nights in the northward direction.

MONTHLY Trends in Clear (Blue) and Usable Time (red) for various sites of astronomical interest










- At ORM there is a larger usable nights fraction in Summer
- In the USA Southwest the effect of the summer monsoon is evident (telescopes shutdown for maintenance)

Cirrus Cloud Parameters Statistics (Base Altitude, Top Altitude, Thickness)








Site	Base Altitude + 1-sigma [km]	Top Altitude + 1-sigma [km]	Cloud Thickness + 1-sigma [meters]
Maunakea	11.7 ± 2.7	12.9 ± 2.7	1257 ± 713
La Palma/ORM	9.3 ± 2.0	10.7 ± 2.1	1332 ± 748
KP/MG	9.6 ± 2.8	10.8 ± 3.0	1188 ± 693
Pachon/Tololo	7.4 ± 3.2	8.2 ± 3.6	885 ± 662
La Silla/Las Campanas	9.5 ± 2.0	10.7 ± 2.1	1152 ± 754
Paranal/Armazones	11.1 ± 2.6	12.1 ± 2.7	1032 ± 602
Cerro Honar	11.7 ± 2.5	12.8 ± 2.5	1068 ± 595

ANNUAL Trends in Optical Depth Cirrus (Blue) and Aerosols (red) for various sites of astronomical interest



- Norther hemisphere sites show a higher level of aerosols than the southern hemisphere sites. (almost a factor of 2)

ANNUAL Trends in Optical Depth Cirrus (Blue) and Aerosols (red) for various sites of astronomical interest



- ORM Site: There is clear evidence of increase in aerosols level (dust) in summer → Calima

Summary of Aerosols & Cirrus Clouds Statistics Maunakea & La Palma only

Maunakea Quartiles	Aerosols optical depth [nepers]	Cirrus clouds optical depth [nepers]	Cirrus Clouds Thickness [m]	Cirrus Clouds Base altitude [km]
25%	0.022	0.015		
50%	0.058	0.056	1257 ± 713	11.7 ± 2.7
75%	0.130	0.130		


La Palma/ORM Quartiles	Aerosols optical depth [nepers]	Cirrus clouds optical depth [nepers]	Cirrus Clouds Thickness [m]	Cirrus Clouds Base altitude [km]
25%	0.029	0.034		
50%	0.067	0.071	1332 ± 748	9.3 ± 2.0
75%	0.146	0.149		

How Cirrus Could Affect AO Performance



- Introducing fluctuations in the laser power propagating from the telescope to the mesosphere.
 - ◇ Therefore reducing the intensity of the LGS photon flux return
- Inducing variations in the intensity of the fratricide effect (in LGS systems with center-launched lasers)
- Inducing photon noise in the WFS sub-apertures that could be harder to calibrate out

Time Variability on Cirrus Optical Depth

$1\sigma \sim 0.1 \text{ mag/airmass (10\%)} \text{ in time scales of 3 secs}$



Cirrus Cloud with max variation $\sim 0.3 \text{ mag/airmass}$



LGS Photon Flux Return

	Optical Depth MK	Optical Depth ORM
Rayleigh	0.0457	0.0576
O3 Chappuis Band	0.0331	0.0331
Aerosols (dust)	0.058	0.067
Cirrus Clouds	0.056	0.071
Total	0.1928	0.2287
Atmospheric Attenuation	0.8246	0.7956

	MK	ORM
Laser Power	20 W	20 W
Zenith Angle	0 deg	0 deg
D2b repumping	10%	10%
Na Atoms (atoms/m ²)	4x10 ¹³	4x10 ¹³
LGS spot size	1.0	1.0
Geomagnetic Field	(that of the site)	(that of the site)
Site Altitude	4050 m	2200 m
LGS Photon return at tel. aperture	14x10⁶ ph/m ² /s	11.3x10⁶ ph/m ² /s

Detection throughput = 0.3477

Frame = 800 Hz

SA size = 0.5² m²

PDE (MK) = 1521 ph/SA/frame

PDE (ORM) = 1228ph/SA/frame


30% increase in cirrus optical depth → 3.6% decrease in PDE

TMT Requirement 900 ph/SA/frame

Zenith Angle Histogram

Keck 2 – one year of pointing


blue-South / red-North




- Ron Holzlohner as shown that in MK the “sweet spot” for LGS return is $\sim 36^\circ$ elevation (54° ZA) which matches the location of the Galactic Center.
- Domenico Bonaccini & collaborators have shown important variability (factor of 2 to 3) in the LGS photon flux return as the laser propagates in directions that benefit less from D2b re-pumping.
- Nightly fluctuations in the Na column density can be large too.
- If we add the effects of cirrus cloud and atmospheric aerosols that will limit the zenith angle range where is possible to meet specifications

TMT Fratricide Pattern on WFS-0

Due to Molecular Rayleigh Backscattered Photons (for Center Launch LGS systems)



Height above ground	Sub Aperture
10 km	4
30 km	11
40 km	14






WFS-0 Aperture Plane
 $SA = \text{int} \{ h * \sin(\theta) / 0.5 \}$

Volume Backscattering Coefficient [m⁻¹ sr⁻¹]

Blue (Maunakea)

Red (ORM)



- At the ORM site, because of its higher density, we expect a ~30% higher level of backscattered photons
- Relative to the ground, Cirrus Clouds will affect rely the same Sub Apertures (independent of the site)

Conclusions

- Within $\pm 5\%$ both sites are comparable regarding the usable number of nights.
- At ORM the cloudiness/precipitation is influenced by the +/- phase of the North Atlantic Oscillation.
- At ORM, there is a clear winter/summer signal in the statistics of usable nights (more in Summer season).
- Chile: a clear trend in the usable nights In the northward direction.
- ORM Site: There is clear evidence of increase in aerosols level (dust) in summer \rightarrow Calima.
- The CALIOP data, seems to indicate the integrated aerosol levels in northern hemisphere is almost a factor of 2 higher than in southern hemisphere sites.
- Cirrus clouds introduce changes in optical depth of order 10%-30% in time scales of few seconds. This affects the laser power able to reach the mesosphere, and the magnitude of the LGS photon flux return. But also, will induce fluctuations in the photons backscattered to the WFS's apertures from the lower levels of the atmosphere.
- Due to the lower altitude (and consequently increase in atmospheric density) it is expected about 30% larger photon noise in the SA affected by fratricide effect.

Conclusions

- Within $\pm 5\%$ both sites are comparable regarding the usable number of nights.
- At ORM the cloudiness/precipitation is influenced by the \pm phase of the North Atlantic Oscillation.
- At ORM, there is a clear winter/summer trend in the number of usable nights (more in Summer season).
- Chile: a clear trend in the number of usable nights.
- ORM Site: There is a clear trend in the number of usable nights (dust) in summer \rightarrow CAI.
- The CAI is a clear trend in the number of usable nights (aerosol levels in northern hemisphere).
- Cirrus clouds are a clear trend in the number of usable nights (aerosol levels in southern hemisphere sites).
- Cirrus clouds are a clear trend in the number of usable nights (depth of order 10%-20% in time scales of order 10-20 years).
- Mesospheric laser power able to reach the mesosphere of the LGS photon flux return. But also, will induce fluctuations in the number of photons backscattered to the WFS's apertures from the lower atmosphere.
- Due to the low altitude (and consequently increase in atmospheric density) it is expected about 30% larger photon noise in the SA affected by fratricide effect.

¡Gracias!

Acknowledgments

The TMT Project gratefully acknowledges the support of the TMT collaborating institutions. They are the Association of Canadian Universities for Research in Astronomy (ACURA), the California Institute of Technology, the University of California, the National Astronomical Observatory of Japan, the National Astronomical Observatories of China and their consortium partners, and the Department of Science and Technology of India and their supported institutes. This work was supported as well by the Gordon and Betty Moore Foundation, the Canada Foundation for Innovation, the Ontario Ministry of Research and Innovation, the National Research Council of Canada, the Natural Sciences and Engineering Research Council of Canada, the British Columbia Knowledge Development Fund, the Association of Universities for Research in Astronomy (AURA), the U.S. National Science Foundation and the National Institutes of Natural Sciences of Japan.

TMT Laser Guide Star Asterisms

3.4.1.1 Asterism generation requirement

[REQ-2-LGSF-0400] The LGSF shall be able to project the NFIRAOS asterism, which consists of 6 LGS, 5 equally spaced on a circle of radius of 35 arcsec and one additional on-axis guide star.

Discussion: NFIRAOS is the TMT early light LGS Multi Conjugate Adaptive Optics (MCAO) system.

[REQ-2-LGSF-0410] The LGSF shall be able to project the MIRA0 asterism, which consists of 3 LGS equally spaced on a circle of radius of 70 arcsec.

[REQ-2-LGSF-0420] The LGSF shall be able to project the MOAO asterism, which consists of 8 LGS, 3 equally spaced on a circle of radius of 70 arcsec and 5 equally spaced on a circle of radius of 150 arcsec.

[REQ-2-LGSF-0430] The LGSF shall be able to project the GLAO asterism, which consists of 5 LGS, 4 equally spaced on a circle of radius of 510 arcsec and one additional on-axis guide star.

[REQ-2-LGSF-0440] The LGSF shall be able to project additional asterisms with up to 9 LGS and radii varying from 5 arcsec to 510 arcsec.

Discussion: The LGSF asterisms for the early light and first decade AO systems are represented in Figure 4.

