

# THE IMPACT OF NON-LTE EFFECTS AND GRANULATION INHOMOGENEITIES ON THE DERIVED IRON AND OXYGEN ABUNDANCES IN METAL-POOR HALO STARS

N. G. SHCHUKINA

Main Astronomical Observatory, National Academy of Sciences, 27 Zabolotnogo Street, Kiev 03680, Ukraine;  
shchukin@mao.kiev.ua

J. TRUJILLO BUENO<sup>1</sup>

Instituto de Astrofísica de Canarias, 38205 La Laguna, Tenerife, Spain; jtb@iac.es

AND

M. ASPLUND

Research School of Astronomy and Astrophysics, Mount Stromlo Observatory, Cotter Road, Weston, ACT 2611, Australia;  
martin@mso.anu.edu.au

Received 2004 May 27; accepted 2004 September 13

## ABSTRACT

This paper presents the results of a detailed theoretical investigation of the impact of non-LTE (NLTE) effects and of granulation inhomogeneities on the derived iron and oxygen abundances in the metal-poor halo subgiant HD 140283. Our analysis is based on both the “classical” one-dimensional stellar atmosphere models and on the new generation of three-dimensional hydrodynamic models. The NLTE calculations presented here have been carried out without inelastic collisions with neutral hydrogen atoms. We find that if NLTE effects are taken into account when synthesizing the Fe I spectrum in both types of atmospheric models, then the derived iron abundance turns out to be very similar in both cases. The emergent spectral line profiles in both models are very much weaker in NLTE than in LTE because the UV overionization mechanism produces a very strong underpopulation of the Fe I levels, in particular in the granular regions of the three-dimensional model. As a result, the NLTE effects on the derived iron abundance are very important, amounting to  $\sim 0.9$  and to  $\sim 0.6$  dex in the three- and one-dimensional cases, respectively. On the other hand, we find that NLTE and three-dimensional effects have to be taken into account for a reliable determination of the iron abundance from weak Fe II lines, because the significant overexcitation of their upper levels in the granular regions tend to produce emission features. As a result, such Fe II lines are weaker than in LTE and the abundance correction amounts to  $\sim 0.4$  dex for the three-dimensional case. We also derive the oxygen-to-iron abundance ratio in the metal-poor star HD 140283 by using the O I triplet at 7772–7775 Å and the forbidden [O I] line at 6300 Å. Our results for the oxygen abundance confirm the values reported in some recent investigations. While the oxygen abundance derived from the O I IR triplet is not very sensitive to the presence of granulation inhomogeneities, such three-dimensional effects amount to  $\sim -0.2$  dex for the [O I] line. The NLTE abundance correction for the O I IR triplet turns out to be  $-0.2$  dex, approximately. Interestingly, when both NLTE and three-dimensional effects are taken into account there still remain significant discrepancies in the iron abundances derived from Fe I and Fe II lines, as well as in the oxygen abundances inferred from the O I and [O I] lines. We conclude that the discrepancies could be due to uncertainties in the stellar parameters of this metal-poor star. We argue that adopting  $T_{\text{eff}} \approx 5600$  K (instead of  $T_{\text{eff}} \approx 5700$  K) and  $[\text{Fe}/\text{H}] \approx -2.0$  (instead of  $[\text{Fe}/\text{H}] \approx -2.5$ ) substantially reduces the discrepancies in the abundances of iron and oxygen inferred from several spectral lines. Under such circumstances, we find  $[\text{O}/\text{Fe}] \approx 0.5$  at  $[\text{Fe}/\text{H}] = -2$ . Obviously, our tentative conclusion that the metallicity of this type of metal-poor star is significantly larger than previously thought may have far-reaching implications in stellar astrophysics.

*Subject headings:* Galaxy: abundances — Galaxy: evolution — nuclear reactions, nucleosynthesis, abundances — stars: atmospheres — stars: late-type — Sun: abundances

## 1. INTRODUCTION

Reliable determinations of the oxygen and iron abundances in the atmospheres of metal-poor stars are of paramount importance for 21st Century astrophysics. This is because the abundances of oxygen and iron in metal-poor stars represent fundamental astrophysical quantities for determining the ages of the oldest stars, for constraining the models of the chemical evolution of the Galaxy, for estimating the yields of light ele-

ments (lithium, beryllium, and boron) from spallation processes in the interstellar and circumstellar gas, for testing different scenarios of nucleosynthesis in supernovae, and so on. Such studies largely rely upon the oxygen-to-iron abundance ratio,  $[\text{O}/\text{Fe}] = \log(\text{O}/\text{Fe})_* - \log(\text{O}/\text{Fe})_{\odot}$ , versus the stellar metallicity  $[\text{Fe}/\text{H}]$ .

### 1.1. The Oxygen Abundance Problem in Metal-Poor Stars

The shape and spread of the  $[\text{O}/\text{Fe}]$  versus  $[\text{Fe}/\text{H}]$  relation is still hotly debated in the literature (e.g., Israelian et al. 2001; Asplund & García Pérez 2001; Meléndez & Barbuy 2002;

<sup>1</sup> Consejo Superior de Investigaciones Científicas; Spain.

Nissen et al. 2002; Fulbright & Johnson 2003; Takeda 2003 and references therein).

The controversy is linked with discordant results obtained using different indicators of the oxygen abundance, such as (1) the oxygen [O I] forbidden  $\lambda 6300$  line, (2) the oxygen O I infrared triplet at 7772–7775 Å, (3) the OH lines in the near ultraviolet, and (4) the OH lines in the infrared. More precisely, the oxygen abundance derived using all these lines agrees more or less with each other for moderate metal-deficient disk stars having  $-1 \leq [\text{Fe}/\text{H}] \leq 0$ . However, there are clear discrepancies among the results obtained for metal-poor halo stars with lower metallicities. Some authors find that the [O/Fe] ratio shows a plateau between 0.4 and 0.5 dex in the metallicity range  $-3 < [\text{Fe}/\text{H}] < -1$  (Barbuy 1988; Kraft et al. 1992; Fulbright & Kraft 1999; Asplund & García Pérez 2001; Carretta et al. 2000; Meléndez & Barbuy 2002), while others find a linear increase with decreasing metallicity, reaching [O/Fe]  $\simeq +1$  dex at  $[\text{Fe}/\text{H}] = -3$  (Abia & Rebolo 1989; Tomkin et al. 1992; Cavallo et al. 1997; Israelian et al. 1998, 2001; Boesgaard et al. 1999; Mishenina et al. 2000 and references therein). Recently, Nissen et al. (2002) have analyzed [O I] and Fe II lines in the framework of three-dimensional atmospheric models and LTE line formation, concluding that there is a quasi-linear trend for  $[\text{Fe}/\text{H}] < -2$ , with [O/Fe]  $\simeq +0.5$  dex at  $[\text{Fe}/\text{H}] = -2.5$ . Furthermore, an extensive one-dimensional reanalysis of published equivalent widths for the O I IR and the forbidden [O I] lines (see Takeda 2003) shows that the discrepancy between O I and [O I] lines in metal-poor halo stars tends to be smaller for stars with higher effective temperatures ( $T_{\text{eff}}$ ) and gravities ( $\log g$ ), i.e., more preferentially for dwarfs and subgiants than for giants. Fulbright & Johnson (2003) support this conclusion. In addition, the oxygen abundances in dwarfs derived from the O I IR lines tend to have values between those obtained with the UV OH and [O I] lines.

### 1.2. The Iron Abundance Problem in Metal-Poor Stars

A key problem directly related with the [O/Fe] issue in metal-poor stars is the reliability of the iron abundance determinations. In fact, the derivation of the iron abundance from Fe I and/or Fe II lines is a very complex problem that requires detailed non-LTE (NLTE) investigations like the one carried by Shchukina & Trujillo Bueno (2001) and in this paper. In general, the Fe I lines suffer from UV overionization in the atmospheres of late-type stars (Athay & Lites 1972; Rutten 1988; Shchukina & Trujillo Bueno 2001). This NLTE effect tends to be substantially more important in metal-poor stars than in solar-like stars as a result of the lower electron density and of the weaker UV blanketing (Thévenin & Idiart 1999; Shchukina & Trujillo Bueno 2001; Israelian et al. 2001, 2004; Shchukina et al. 2003). We should mention also that Gratton et al. (1999) concluded that NLTE effects are rather small in metal-poor stars. However, their calculations presumably overestimated the efficiency of inelastic collisions with hydrogen atoms, which thermalize the Fe I levels and thus almost restore LTE conditions.

It is of interest to point out that while the Fe I lines are (in principle) expected to be sensitive to the UV overionization mechanism, the LTE approximation is thought to be suitable for weak lines of Fe II (see, e.g., Asplund et al. 1999; Nissen et al. 2002). However, as we shall see below, this does not necessarily have to be the case, because many subordinate Fe II lines whose upper levels are of odd parity (starting at  $z^6D^o$ ) might be partially filled by emission owing to an optical pumping mecha-

nism similar to that investigated by Cram et al. (1980) for the solar case.

### 1.3. The Solar Oxygen Abundance Problem

Obviously, the solar case constitutes a fundamental reference for a reliable quantification of chemical abundances in other stars. The solar iron abundance appears to be fairly well determined now [ $\log \epsilon_{\odot}(\text{Fe}) = 7.50 \pm 0.10$ ; see Shchukina & Trujillo Bueno 2001]—that is, it is similar to the meteoritic value. In fact, this is the value adopted by Nissen et al. (2002) in their study of the O/Fe ratio in metal-poor stars.

Great progress has been also made toward a reliable determination of the solar oxygen abundance. The one-dimensional modeling approach leads to discrepancies between the solar oxygen abundance determined from the permitted O I lines and from the [O I] and OH lines. Abundance determinations from OH lines using the one-dimensional model of Holweger & Müller (1974) suggested first a high oxygen abundance value [ $\log \epsilon_{\odot}(\text{O}) = 8.93 \pm 0.03$ ; see Grevesse et al. 1984; Sauval et al. 1984; Anders & Grevesse 1989]. Several years later, Grevesse & Sauval (1998) obtained an oxygen abundance that was lower by 0.1 dex. This lower value was found after using improved  $\log gf$  values and a slightly modified one-dimensional model, which helped to remove the existence of a dependency of the derived oxygen abundance with the excitation potential of the lower level of the OH transitions. The forbidden [O I] line gives also a high oxygen abundance value when the contribution of the blend due to a Ni I line is ignored (Lambert, 1978; Sauval et al. 1984; Holweger 2001). However, the permitted O I lines at 7772–7775 Å yield a value for the oxygen abundance that is lower by 0.2 dex (Kiselman 1993; Holweger 2001). Some authors have argued that the existing NLTE calculations for these spectroscopic indicators of the oxygen abundance might not be valid owing to the neglect of inelastic collisions with hydrogen. For this reason, the solar oxygen abundance derived from the [O I] forbidden line was considered to be the most reliable one, because of the suitability of the LTE approximation (hence, uncertainties of collisional rates with hydrogen atoms do not play any role). Another argument used to support a high value for the oxygen abundance is the small scatter in the oxygen abundances obtained from different OH lines. Kiselman & Nordlund (1995) tried to clarify the solar oxygen abundance issue by using a three-dimensional solar photospheric model that is similar to, but less sophisticated than, that of Asplund et al. (2000). They concluded that a downward revision of the solar oxygen abundance is required, with respect to the value derived from [O I] and OH lines. More recently, Allende Prieto et al. (2001) have obtained a more precise estimate by taking into account the influence of the above-mentioned blend by the Ni I line and by using the three-dimensional hydrodynamic model of Asplund et al. (2000). They found  $\log \epsilon_{\odot}(\text{O}) = 8.69 \pm 0.05$ , which shows that the [O I]–based abundance derived via the three-dimensional approach is significantly lower than that derived from classical one-dimensional model atmospheres.

At a recent IAU symposium on modeling of stellar atmospheres, Shchukina et al. (2003) reported that the oxygen abundance obtained via NLTE synthesis of the solar O I IR triplet in the three-dimensional solar photosphere model of Asplund et al. (2000) is  $\log \epsilon_{\odot}(\text{O}) \approx 8.7$ . This result is in good agreement both with the conclusion of Allende Prieto et al. (2001) and with the recent three-dimensional analysis of Asplund et al. (2004), who found excellent agreement between

the averaged oxygen abundance values derived from [O I], O I, OH vibration-rotation, and OH pure rotation lines.<sup>2</sup>

#### 1.4. Possible Reasons for the Abundance Discrepancies

It is currently believed that abundance discrepancies in metal-poor halo stars and the Sun are dominated by deficiencies in the spectral line modeling. Errors in the measurements of equivalent widths are of minor importance, with the exception of weak [O I] and Fe II lines. All abundance indicators in metal-poor stars are to a different degree sensitive to the adopted stellar parameters ( $T_{\text{eff}}$ ,  $\log g$ , and [Fe/H]). They have to depend also on atmospheric properties such as the temperature gradient and the ratio of the line to continuum opacity. Large uncertainties could result from neglecting atmospheric inhomogeneities and velocity fields (i.e., granulation inhomogeneities produced by the stellar surface convection process). Part of the discrepancies are probably due to NLTE effects. There is now an abundance of literature concerning the iron and oxygen line formation problems in classical plane-parallel stellar atmospheric models (Thévenin & Idiart 1999; Mishenina et al. 2000; Kiselman 2001; Takeda 2003 and references therein). The recent generation of three-dimensional hydrodynamic models of stellar atmospheres have activated the present debates on the influence of granulation inhomogeneities and of NLTE effects on the determination of chemical abundances in metal-poor stars (see Asplund et al. 1999, 2004; Asplund & García Pérez et al. 2001; Shchukina & Trujillo Bueno 2001; Nissen et al. 2002). In this respect, it is very important to emphasize that the three-dimensional hydrodynamic simulations of Asplund and co-workers show that the atmospheres of metal-poor halo stars may have substantially lower kinetic temperatures than the radiative equilibrium temperatures of the “classical” theoretical one-dimensional model atmospheres of Kurucz (1993) and the MARCS collaboration (Gustafsson et al. 1975).

#### 1.5. Motivation

The previous discussion makes evident that detailed, 3D+NLTE investigations of the iron and oxygen abundances are urgently needed. However, it is crucial that this type of analyses be carried out *jointly* for iron and oxygen, and using the same set of atmospheric models, continuum opacity package, and radiative transfer code. The present paper presents the results of such an investigation. We study the NLTE formation problem of the Fe I, Fe II, and O I lines in a three-dimensional hydrodynamic model of the atmosphere of the metal-poor star HD 140283. This classical halo subgiant is widely used for studies of the very early Galaxy and the origin of the chemical elements. In order to quantify the impact of three-dimensional simulations on the abundance determinations in HD 140283, we carry out a detailed comparison with the one-dimensional classical modeling. To this end, we use the three-dimensional hydrodynamic atmospheric model of Asplund’s et al. (1999) for the star HD 140283, as well as a suitable grid of Kurucz’s one-dimensional models. Moreover, we evaluate the extent to which 3D+NLTE effects influence the determination of the stellar parameters of this metal-poor subgiant.

<sup>2</sup> It is interesting to mention that the oxygen abundances obtained by Asplund et al. (2004) from OH rotational lines using the three-dimensional modeling approach show a large line-to-line scatter, and with sensitivity to the line strength and to the excitation potential of the transition’s lower level. However, such presumably minor discrepancies are not the topic of this paper. Here we restrict ourselves to consider the NLTE determination of the solar oxygen abundance via O I lines, for consistency with similar computations that we have carried out for the metal-poor star HD 140283.

Our paper considers also the case of the Sun as a reference for stellar abundance determinations with emphasis on the solar oxygen abundance derived from the O I IR triplet. As in our 3D+NLTE analysis of the solar iron abundance (Shchukina & Trujillo Bueno 2001) we investigate here the solar oxygen abundance problem by using the same three-dimensional hydrodynamic solar photosphere model.

## 2. THE DIAGNOSTIC METHOD

In order to determine the stellar abundances of iron and oxygen, we have to compare some of the properties of the observed spectral line profiles (e.g., their equivalent widths) with those calculated via the numerical solution of the radiative transfer equation in a given model of the atmosphere of the star under investigation. To this end, we have used the same NLTE code and strategy that we applied in our previous investigation of the iron line formation problem in a three-dimensional hydrodynamic model of the solar photosphere (see Shchukina & Trujillo Bueno 2001), with the only difference that we have now computed flux profiles. Our three-dimensional strategy for investigating the issue of the abundances of iron and oxygen in the metal-poor star HD 140283 is similar to that of Shchukina & Trujillo Bueno (2001) for solar-like atmospheres, but using a three-dimensional snapshot model from the Asplund et al. (1999) stellar surface convection simulation. Therefore, we have neglected the effects of horizontal radiative transfer on the atomic level populations. As discussed by Shchukina & Trujillo Bueno (2001), this is expected to be a good approximation for determining the stellar iron abundance from Fe I and Fe II lines. Kiselman & Nordlund (1995) already demonstrated that it is also a good approximation for the oxygen IR triplet.

Our NLTE code is based on very efficient iterative methods (Trujillo Bueno & Fabiani Bendicho 1995; Socas-Navarro & Trujillo Bueno 1997; Trujillo Bueno 2003) that were developed to facilitate radiative transfer simulations with complex atomic systems in realistic stellar atmospheric models. We have carried out both LTE and NLTE spectral line syntheses by using both the one- and three-dimensional models mentioned below. In each case, the abundances of iron and oxygen are modified iteratively until the best fit to the observations is achieved.

## 3. THE OBSERVATIONS

We have chosen 33 Fe I and 15 Fe II lines for our three- and one-dimensional determinations of the iron abundance in the halo subgiant HD 140283. The equivalent widths ( $W$ ) of all the Fe I lines and of 11 Fe II lines were obtained from high-resolution spectra observed a few years ago with the 2.7 m Harlan J. Smith Telescope at McDonald observatory (R. García López 2000, private communication; see also Allende Prieto et al. 1999). The equivalent widths ( $W$ ) for the remaining four Fe II lines are taken from Table 3 of Nissen et al. (2002). They are weak lines with  $W$  in the range 1–4 mÅ. Their wavelengths are  $\lambda\lambda 6149.238$ , 6247.562, 6432.654, and 6456.376. The iron lines used in our analysis are listed in Table 1.

The O I IR triplet lines in the metal-poor star HD 140283 are weak. Given uncertainties in the measurements, we have averaged their equivalent widths over three sets of observations (Abia & Rebolo 1989; Tomkin et al. 1992; G. Israelian 2003, private communication). The resulting equivalent widths of the  $\lambda\lambda 7772$ , 7774, and 7775 lines used in our oxygen abundance determinations are 7.9, 4.8, and 3.4 mÅ, respectively. The equivalent width of the forbidden [O I]  $\lambda 6300$  line ( $W = 0.5$  mÅ)

TABLE 1  
O I, Fe I AND Fe II LINE LIST AND OBSERVED EQUIVALENT WIDTHS ( $W$ )  
FROM THE METAL-POOR STAR HD 140283

Ion	$\lambda$ (Å)	$\log gf$	$\chi$ (eV)	$W$ (mÅ)
O I	6300.230	-9.750	0.00	0.5
O I	7771.960	0.324	9.11	7.9
O I	7774.180	0.174	9.11	4.8
O I	7775.400	-0.046	9.11	3.4
Fe I	3906.482	-2.243	0.11	78.5
Fe I	3917.185	-2.155	0.99	44.1
Fe I	4005.246	-0.610	1.55	79.4
Fe I	4147.673	-2.104	1.48	24.1
Fe I	4152.172	-3.232	0.95	9.1
Fe I	4177.597	-3.058	0.91	24.3
Fe I	4202.031	-0.708	1.48	78.3
Fe I	4216.186	-3.356	0.00	38.9
Fe I	4222.219	-0.967	2.44	30.5
Fe I	4250.125	-0.405	2.46	52.4
Fe I	4271.159	-0.349	2.44	60.0
Fe I	4375.932	-3.031	0.00	55.0
Fe I	4447.722	-1.342	2.21	25.0
Fe I	4489.741	-3.966	0.12	11.2
Fe I	4494.568	-1.136	2.19	34.8
Fe I	4733.596	-2.970	1.48	5.5
Fe I	4939.690	-3.340	0.86	9.9
Fe I	4994.133	-3.080	0.91	13.8
Fe I	5012.071	-2.642	0.86	32.3
Fe I	5083.342	-2.958	0.95	15.5
Fe I	5110.414	-3.760	0.00	24.8
Fe I	5123.723	-3.068	1.01	12.3
Fe I	5194.943	-2.090	1.55	25.7
Fe I	5198.714	-2.135	2.21	8.5
Fe I	5429.699	-1.879	0.95	67.9
Fe I	5701.553	-2.216	2.55	2.5
Fe I	6065.487	-1.530	2.60	9.9
Fe I	6136.620	-1.400	2.44	19.7
Fe I	6219.290	-2.433	2.19	3.2
Fe I	6252.561	-1.687	2.39	14.4
Fe I	6265.140	-2.550	2.17	5.0
Fe I	6593.878	-2.422	2.42	5.5
Fe I	6750.152	-2.621	2.41	3.1
Fe II	4173.450	-2.180	2.57	27.1
Fe II	4178.855	-2.480	2.57	20.3
Fe II	4303.166	-2.490	2.69	19.1
Fe II	4416.817	-2.600	2.77	11.2
Fe II	4491.401	-2.700	2.84	7.9
Fe II	4555.890	-2.290	2.82	16.9
Fe II	4583.829	-2.020	2.79	41.5
Fe II	4923.921	-1.320	2.88	57.2
Fe II	5197.569	-2.380	3.22	10.2
Fe II	5234.620	-2.310	3.21	13.2
Fe II	5316.609	-1.850	3.14	24.1
Fe II	6149.238	-2.850	3.87	1.2
Fe II	6247.562	-2.510	3.87	2.1
Fe II	6432.654	-3.740	2.88	1.3
Fe II	6456.376	-2.280	3.89	4.3

NOTES.—The oscillator strengths for the O I IR triplet are from Kurucz & Bell (1995), while that for the forbidden [O I] line is from Lambert (1978). The wavelengths and excitation energies of the Fe lines are from Moore (1959). The Fe oscillator strengths were taken from Fuhr et al. (1988), except for Fe I  $\lambda\lambda$ 4733.596 and Fe II  $\lambda\lambda$ 5197.569, 5234.620, 6149.238, and 6456.376, which stem from Gurtovenko & Kostik (1989). The observed equivalent widths of all the Fe I lines and of 11 Fe II lines have been kindly provided to us by R. García López (2000, private communication), while the equivalent widths of the remaining Fe II lines have been taken from Table 3 of Nissen et al. (2002).

has been taken from Nissen’s et al. (2002) paper. The oxygen line data are specified in Table 1.

Measurements of the equivalent widths of the solar O I IR lines tend to have an uncertainty of  $\sim 5\%$ . Therefore, we have used averaged values of five sets of observations at the solar disk center (Delbouille et al. 1973; Gurtovenko & Kostik 1989; Altrock 1968; Müller et al. 1968; King & Boesgaard 1995). The resulting averaged solar equivalent widths for the  $\lambda\lambda$ 7771.96, 7774.18, and 7775.40 lines are 81.6, 70.4, and 57.2 mÅ, respectively.

#### 4. THE ATMOSPHERIC MODELS

In each case (i.e., the solar and the metal-poor star), we have used a single three-dimensional snapshot model taken from realistic radiation hydrodynamic simulations of stellar surface convection (Stein & Nordlund 1998; Asplund et al. 1999, 2000). Such time-dependent simulations have no adjustable free parameters, since they are based on the numerical solution of the equations of mass, momentum, and energy conservation for given values of the effective temperature ( $T_{\text{eff}}$ ), surface gravitational acceleration ( $\log g$ ), and chemical composition. In order to facilitate the radiative transfer calculations, the original snapshots, which have 82 grid points along the vertical direction, were interpolated to grids of  $50 \times 50 \times 102$  and  $50 \times 50 \times 121$  points for the solar and metal-poor star cases, respectively. The improvement in the vertical resolution (i.e., with 102 and 121 grid points sampling the line and continuum forming regions) was introduced to enhance the accuracy of the NLTE calculations.

The solar three-dimensional model is that described by Asplund et al. (2000). This three-dimensional model of the solar photosphere has been used by Shchukina & Trujillo Bueno (2001) in their NLTE investigation of the solar iron abundance. We have also used it to determine the solar oxygen abundance via NLTE analysis of the O I IR triplet at 7772–7775 Å. Information on the corresponding three-dimensional model for the star HD 140283 can be found in Asplund et al. (1999). The adopted stellar parameters were  $\log g = 3.7$ ,  $T_{\text{eff}} = 5700$  K, and  $[\text{Fe}/\text{H}] = -2.5$ , which result from the infrared flux method (IRFM) for  $T_{\text{eff}}$ , from *Hipparcos* parallaxes for  $\log g$  and from published values for the  $[\text{Fe}/\text{H}]$  (see Asplund et al. 1999). In our differential comparison between the results obtained via the one- and three-dimensional approaches we have used atmospheric models characterized by the same stellar parameters ( $T_{\text{eff}}$ ,  $\log g$ , and  $[\text{Fe}/\text{H}]$ ). In § 8 we will also show results in a grid of “classical” one-dimensional models spanning the ranges  $5300 \leq T_{\text{eff}} \leq 5700$ ,  $3.7 \leq \log g \leq 3.9$ , and  $-2.0 \leq [\text{Fe}/\text{H}] = -2.5$ .

#### 5. THE ATOMIC MODELS

In our spectral synthesis of oxygen and iron lines we adopted the quantum-mechanical approach developed by Anstee & O’Mara (1995), Barklem & O’Mara (1997), and Barklem et al. (1998) for the van der Waals collisional broadening. In the following subsections we provide some information about the atomic models of oxygen and iron we have used for our radiative transfer calculations.

##### 5.1. Oxygen

Our atomic model for oxygen is based on the data of Carlsson & Judge (1993). It has 23 O I fine-structure levels and one O II level. We considered 31 bound-bound and 23 bound-free radiative transitions. The O I Grotrian diagram has a helium-like

TABLE 2  
SOLAR OXYGEN AND IRON ABUNDANCES OBTAINED BY DIFFERENT AUTHORS USING THE ASPLUND ET AL. (2000) THREE-DIMENSIONAL HYDRODYNAMIC MODEL ATMOSPHERE AND THE ONE-DIMENSIONAL PHOTOSPHERIC MODEL OF HOLWEGER & MÜLLER (1974)

Lines	Approach	Present Study	Asplund et al. (2004)	Allende Prieto et al. (2001)	Shchukina & Trujillo Bueno (2001)	Kostik et al. (1996)
Three-dimensional Solar Model						
O I IR.....	NLTE	8.70 ± 0.06	8.65 ± 0.01	...	...	...
O I IR.....	LTE	8.93 ± 0.06	8.89 ± 0.02	...	...	...
[O I].....	LTE	...	8.68 ± 0.01	8.69	...	...
OH vibration-rotation.....	LTE	...	8.61 ± 0.03	...	...	...
OH rotation.....	LTE	...	8.65 ± 0.02	...	...	...
Fe I.....	NLTE	...	...	...	7.50 ± 0.10	...
Fe I.....	LTE	...	...	...	7.43 ± 0.11	...
One-dimensional Model						
O I IR.....	NLTE	8.64 ± 0.06	8.61 ± 0.01	...	...	...
Fe I.....	LTE	...	...	...	...	7.62 ± 0.04

NOTES.—The given uncertainties reflect only the line-to-line scatter. The results are based on permitted O I IR lines, forbidden [O I] lines, OH vibration-rotation lines, OH pure rotation lines and lines of Fe I.

structure. The triplets and the quintets are only weakly coupled to each other. The two systems differ in one important aspect: the triplets are connected to the ground state by strong resonance lines, while the quintets are not. The latter form a separate system with the  $3^5S$  level acting as a ground level and the  $\lambda\lambda 7772-7775$  IR triplet as resonance lines. O I levels have hydrogen-like energies. As a result, there is an important wavelength coincidence of the O I  $\lambda 1025$  line with the hydrogen Ly $\beta$  line. We emphasize that the NLTE results for O I are quite insensitive to including additional levels and transitions (Shchukina 1987; Kiselman 1993; Takeda 2003). In agreement with previously published results, we have found that the effects of the coupling between the triplet and quintet systems, Ly $\beta$  pumping, and the binding of CO molecules are marginal in the stellar atmospheres studied here (see Shchukina 1987; Kiselman 1993; Carlsson & Judge 1993; Kiselman & Nordlund 1995).

### 5.2. Iron

Our model atom for Fe I+Fe II+Fe III is similar to that used by Shchukina & Trujillo Bueno (2001). It has 225 Fe I levels, 23 Fe II levels, and one Fe III level, including their multiplet fine structure. The Fe I levels are interconnected by 330 bound-bound radiative transitions, while we have 25 strong radiative transitions among the Fe II levels. All levels are coupled via collisional transitions with electrons. The Fe I term diagram is, in fact, complete up to an excitation potential  $\chi = 5.72$  eV. At higher energies it contains about 50% of the terms that have been presently identified. Each term of our atomic model is coupled to a parent term of the next ionization stage by photoionization transitions. For most of the Fe I terms that have  $\chi \leq 5$  eV, we used the experimental photoionization cross sections given by Athay & Lites (1972). For the rest of them, and also for the Fe II terms, the hydrogen-like approximation was used.<sup>3</sup>

In our calculations we have not included inelastic collisions with hydrogen atoms. We have also neglected the UV haze opacities. We have carried out several numerical experiments in

<sup>3</sup> It is interesting to mention that the new radiative cross sections of Bautista (1997) are substantially larger than those given by the hydrogenic approximation. Therefore, in respect to the influence of photoionization cross sections, we might perhaps be underestimating the NLTE effects.

order to investigate how large the uncertainties in the iron abundance determinations could be as a result of uncertainties in the hydrogen collisional rates, in the UV haze opacity, and in the bound-free cross sections. Considering all factors together, we have found that the uncertainty for the solar iron abundance does not exceed 0.07 dex (see Shchukina & Trujillo Bueno 2001). The problem of the iron abundance uncertainty for the metal-poor star HD 140283 is discussed in § 8.

## 6. THE OXYGEN ABUNDANCES

In this section we report on the results we have obtained for the oxygen abundance in the Sun and in the metal-poor star HD 140283. Our abundance determinations and some recent abundance studies by other authors are briefly summarized in Tables 2 and 3.

### 6.1. The Sun

We have derived the abundance of oxygen in the solar photosphere by fitting the observed disk center profiles and equivalent widths of the O I IR triplet via spectral synthesis in the three-dimensional model of the solar photosphere. Taking into account NLTE effects, we find  $\log \epsilon(\text{O}) = 8.70 \pm 0.06$  (see also Shchukina et al. 2003), while the LTE approximation gives  $\log \epsilon(\text{O}) = 8.93 \pm 0.06$ . Note that the uncertainties reflect the line-to-line scatter for the O I IR triplet lines. In both cases we obtained an almost perfect fit to the observed disk center profiles and equivalent widths. Our NLTE result is in good agreement with that obtained by Allende Prieto et al. (2001) from the forbidden [O I]  $\lambda 6300$  line [ $\log \epsilon(\text{O}) = 8.69$ ]. It is also consistent with the three-dimensional result of Asplund et al. (2004) for the OH lines [ $\log \epsilon(\text{O}) = 8.61 - 8.65$ ] and with their full 3D+NLTE result for the O I triplet [ $\log \epsilon(\text{O}) = 8.65$ ]. It is important to note that the small discrepancy between our solar abundance value derived from the O I IR triplet and that of Asplund et al. (2004) is mainly due to differences in the oscillator strengths used ( $\Delta \log gf \approx -0.05$  dex). The oxygen abundance changes that result from a 5% uncertainty in the solar equivalent width measurements of these spectral lines lie in the same range ( $\approx \pm 0.05$  dex).

The NLTE formation of the O I triplet can be easily understood via a two-level atom approach if the line opacity is calculated assuming LTE (Shchukina 1987; Kiselman 1993;

TABLE 3  
OXYGEN AND IRON ABUNDANCES FOR HD 140283 OBTAINED BY DIFFERENT AUTHORS USING THE ASPLUND ET AL. (1999) THREE-DIMENSIONAL HYDRODYNAMIC MODEL ATMOSPHERE AND ONE-DIMENSIONAL HYDROSTATIC MODELS

Lines	Approach	Present study	Nissen et al. (2002)	Asplund et al. (1999)
Three-dimensional Model				
O I IR.....	NLTE	7.08 ± 0.06	...	...
O I IR.....	LTE	7.24 ± 0.06	...	7.20 ( $\lambda 7772$ )
[O I].....	LTE	6.96	6.83	...
Fe I.....	NLTE	5.77 ± 0.19	...	...
Fe I.....	LTE	4.85 ± 0.15	...	4.57 ± 0.16
Fe II.....	NLTE	5.60 ± 0.48	...	...
Fe II.....	LTE	5.18 ± 0.11	5.13 ± 0.06	5.16 ± 0.10
One-dimensional Models				
O I IR.....	NLTE	7.11 ± 0.06	7.02	...
O I IR.....	LTE	7.29 ± 0.06	7.11	7.15 ( $\lambda 7772$ )
[O I].....	LTE	7.18	7.09	...
Fe I.....	NLTE	5.77 ± 0.15	...	...
Fe I.....	LTE	5.20 ± 0.14	...	5.02 ± 0.17
Fe II.....	NLTE	5.34 ± 0.21	...	...
Fe II.....	LTE	5.18 ± 0.11	5.08 ± 0.06	5.08 ± 0.11

NOTES.—The given uncertainties reflect the line-to-line scatter. In general, the abundances are based on different selection of lines,  $gf$ -values and/or equivalent widths. The results given in the table are based on permitted O I IR lines, forbidden [O I] lines, and Fe I and Fe II lines. Parameters for all models are  $T_{\text{eff}} = 5700$ ,  $\log g = 3.7$ , and  $[\text{Fe}/\text{H}] = -2.5$ , except for the one-dimensional model of Nissen et al. (2002), which are  $T_{\text{eff}} = 5690$ ,  $\log g = 3.69$ , and  $[\text{Fe}/\text{H}] = -2.42$ .

Kiselman & Nordlund 1995; Takeda 2003; Shchukina et al. 2003; Asplund et al. 2004 and references therein). The photon losses mechanism lowers the line source functions both in the granular and intergranular regions of the three-dimensional model, which implies that the NLTE profiles of the IR triplet lines are deeper than in LTE. The effect is larger in the intergranular regions, but the spatially averaged emergent profiles are only slightly affected by the granulation inhomogeneities. Consequently, the effects of granulation inhomogeneities on the NLTE solar oxygen abundance determination are small, i.e.  $\sim 0.06$  dex (Shchukina et al. 2003). More significant are the NLTE abundance corrections for the O I infrared triplet lines observed at the solar disk center, which amount to  $-0.25$ ,  $-0.22$ , and  $-0.20$  dex for  $\lambda\lambda 7771.96$ ,  $7774.18$ , and  $7775.40$ , respectively—that is, the LTE approximation tends to overestimate the derived oxygen abundance. Such NLTE effects are very similar to those reported by Asplund et al. (2004), which result from full 3D+NLTE computations.

### 6.2. The Metal-Poor Halo Subgiant HD 140283

It is convenient to begin this section by giving the values of the oxygen abundances we have obtained *without* assuming LTE when using the one- and three-dimensional models of this metal-poor star. As mentioned above, both models are characterized by  $\log g = 3.7$ ,  $T_{\text{eff}} = 5700$  K, and  $[\text{Fe}/\text{H}] = -2.5$ . The one-dimensional model gives  $\log \epsilon(\text{O}) = 7.11 \pm 0.06$  for the O I IR triplet and  $\log \epsilon(\text{O}) = 7.18$  for the [O I] forbidden line at 6300 Å. Concerning the three-dimensional analysis, we find  $\log \epsilon(\text{O}) = 7.08 \pm 0.06$  when using the O I IR triplet but  $\log \epsilon(\text{O}) = 6.96$  for the [O I] line. Our one-dimensional results differ slightly from those presented by Nissen et al. (2002), probably because we have not used the same continuum opacity package and/or identical one-dimensional atmospheric models. Our estimates indicate that the abundance differences could reach  $\sim 0.05$ – $0.07$  dex.

Having given these values, it is now of interest to note that the NLTE effects in the one- and three-dimensional atmospheric models of this metal-poor star are not negligible, but they are smaller than in the solar case because the O I IR lines originate in deeper layers where their source functions and the level populations of their upper levels are closer to their LTE values (Shchukina et al. 2003). Adopting LTE for the O I lines leads to an abundance overestimation of 0.18 and 0.16 dex for the one- and three-dimensional cases, respectively. The determination of the oxygen abundance based on the O I IR triplet is nearly insensitive to three-dimensional effects. The correction does not exceed 0.03 dex.

On the contrary, the forbidden [O I] line at 6300 Å is sensitive to the stellar atmospheric inhomogeneities. In agreement with Nissen et al. (2002), we find that a three-dimensional analysis reduces the oxygen abundance derived from this forbidden line by 0.22 dex. The NLTE correction is extremely small ( $< -0.01$  dex).

## 7. THE IRON ABUNDANCE

This section focusses on the iron abundance in the Sun and in HD 140283. Our results and a comparison with those of recent publications are given in Tables 2 and 3.

### 7.1. The Sun

Shchukina & Trujillo Bueno (2001) applied a 3D+NLTE approach to determine the solar iron abundance from 18 Fe I lines that were carefully selected. Ideally, one would like to determine the iron abundance in the Sun and HD 140283 employing the same Fe I line list. However, because of the relatively high solar metallicity the solar spectral lines at wavelengths below 5000 Å are much more affected by blends than in the metal-poor star HD 140283. As a result, overlapping wings of nearby strong lines cause large uncertainties in the equivalent width measurements of most of the solar Fe I lines listed in Table 1. The only




FIG. 1.—Departure coefficients ( $\beta$ ) vs. height in the three-dimensional atmospheric model of the metal-poor star HD 140283. *Left*: Representative “granular” point. *Right*: Representative “intergranular” point. The solid lines indicate the  $\beta$ -coefficients for the Fe I levels with excitation potentials below 5 eV. The dashed and dash-dotted lines refer to the  $\beta$ -coefficients of the lower and upper levels of the Fe II lines in Table 1, respectively. The thick lines correspond to the Fe I ground level  $a^5D$ , to the Fe II ground level  $a^6D$ , and to the Fe II-excited level of odd parity  $z^6D^o$ , respectively. Arrows mark the representative atmospheric heights below which one of the Fe I lines ( $\lambda 5429.699$ ) and the Fe II lines from Table 1 originate in the NLTE and LTE cases. Note that such NLTE and LTE heights are nearly the same for the Fe II lines.

exceptions are the lines at  $\lambda > 5700 \text{ \AA}$  (i.e., only eight of 33 Fe I lines). In any case, the main reason that the same lines cannot be used in the Sun and HD 140283 is that the line strengths are vastly different in the two cases. The weak iron lines in HD 140283 are very strong and therefore ill-suited for abundance purposes in the Sun. Thus, to recalculate the iron abundance of the Sun using the lines of Table 1 is hardly possible.

The solar Fe I lines used in our three-dimensional solar abundance determination (see Table 1 in Shchukina & Trujillo Bueno 2001) were selected from the well-known “Kiel+Oxford” line list according to the criteria discussed by Kostik et al. 1996. The lines chosen by Shchukina & Trujillo Bueno (2001) account for the iron abundance behavior versus the lower level excitation potential  $\chi$ —that is, for the existence of both an abundance dependence on  $\chi$  and a sizable scatter in the abundance values for each  $\chi$ . These 18 representative Fe I lines yield the largest and lowest abundances for each  $\chi$ -value and span a large range in  $\chi$  (0–4.6 eV). This should be a very suitable choice of lines for evaluating the largest three-dimensional effects. Shchukina & Trujillo Bueno (2001) obtained an iron abundance  $\log \epsilon_{\odot}(\text{Fe}) = 7.50 \pm 0.10$  for the 3D+NLTE case. The 3D+LTE approach gives a lower abundance, i.e.,  $\log \epsilon_{\odot}(\text{Fe}) = 7.43 \pm 0.11$ . Neglecting the effect of granulation inhomogeneities leads to a considerably larger value [ $\log \epsilon_{\odot}(\text{Fe}) = 7.62 \pm 0.04$ ; e.g. Kostik et al. 1996].

We stress that the Fe I lines we have used for the star HD 140283 were chosen to obtain a combination of lines that is comparable to that provided by the above-mentioned 18 solar lines (with respect to their equivalent widths and  $\chi$  coverage).

### 7.2. The Metal-Poor Halo Subgiant HD 140283

Like in the solar case (Shchukina & Trujillo Bueno 2001), near-UV overionization appears to be the main NLTE mechanism for Fe I in this metal-poor star. In both the one- and three-dimensional models it causes an important underpopulation of the Fe I levels and hence a reduction in the opacity of the Fe I lines (Fig. 1). As a result, the atmospheric heights where the line optical depth reaches unity are shifted toward deeper layers where the line source functions turn out to be close to the Planck

function. The relevance of this effect increases with the vertical temperature gradient, which is larger above the granules. For this reason, we find a strong weakening of the Fe I lines that emerge from such granular regions, while above the intergranular lanes the emergent spectral line profiles do not change appreciably. On the whole, the calculated NLTE profiles turn out to be much weaker than when the LTE approximation is used (see Fig. 2).

Concerning NLTE effects in Fe II lines, it is very important to point out that in the granular regions we find a significant overpopulation of the excited levels of odd parity starting at  $z^6D^o$  (Fig. 1, right). According to Cram et al. (1980) and Rutten (1988), this effect is caused by an excess in the UV continuum radiation field near  $2600 \text{ \AA}$ , which leads to a pumping of the upper level populations of the Fe II lines. As a consequence, the source functions of the Fe II lines with high excitation energy of the upper level ( $\chi_{\text{up}} \geq 4.8 \text{ eV}$ ) exceed significantly the Planck function values. This implies that such Fe II lines are weaker than stipulated by the LTE approximation and may even be in emission instead of in absorption. This radiative pumping mechanism takes place in both the one- and three-dimensional models. Like in the case of Fe I lines, it is sensitive to the vertical temperature gradient. The steeper the gradient, the larger the pumping. As a result, in the granular regions where the temperature gradients are larger the effect is much more pronounced (see Fig. 1).

The importance of such an effect is illustrated in Figure 3, which shows both NLTE and LTE results for two Fe II lines using the three-dimensional model atmosphere of the metal-poor subgiant HD 140283. We point out that emission is much more pronounced for the emergent profiles that originate above the stellar granular regions, while above the intergranular lanes the spectral line profiles are hardly modified with respect to the LTE case. Allowing for departures from LTE produces a drastic change in the calculated spatially averaged flux of the weak Fe II  $\lambda 6247.562$  line in the three-dimensional model: the absorption profile is replaced by one in emission. Note that this spectral line is taken from the set of weak Fe II lines that Nissen et al. (2002) assumed to be insensitive to NLTE effects. However, we




FIG. 2.—Emergent disk-center intensity profiles of the Fe I  $\lambda 5429.699$  line calculated in the three-dimensional hydrodynamic model of the metal-poor star HD 140283. The left panels correspond to the NLTE case, while the right panels to LTE. *Top panels:* The wavelength variation of the emergent intensity that one would be able to extract at each point along the spectrograph’s slit if the “surface” of the star HD 140283 could be spatially resolved. The “wiggles” are caused by the Doppler shifts associated with the velocity flows of the stellar granulation pattern. Bright strips correspond to granules, while dark ones corresponded to intergranules. The hypothetical slit is located along the position  $Y = 24$  in the three-dimensional model. Intensities are given in absolute energy units ( $\text{ergs cm}^{-2} \text{s}^{-1} \text{sr}^{-1} \text{Hz}^{-1}$ ). Black lines show variations of the line equivalent widths  $W$  along the slit. *Bottom panels:* The spatially resolved emergent intensity profiles at each of the 50 surface grid points considered. The thick solid lines with filled circles show the resulting spatially averaged intensity.

find that this type of lines do suffer from the above-mentioned UV pumping mechanism because all of them belong to the aforementioned category of lines, i.e., their upper levels are of odd parity and are situated above the  $z^6D^o$  level. The stronger Fe II  $\lambda 5316.609$  line displayed in Fig. 3 is also from this category. Figure 3 clearly demonstrates that the stronger the spectral line the smaller the relative effect of the UV-pumping. For this line emission arises only at the core of the “granular” profiles. Spatial averaging completely masks such an emission but makes the flux profile weaker than in the LTE. It is also of interest to mention that in the one-dimensional stellar atmosphere model departures from LTE produce similar changes in the Fe I and Fe II lines, but they are less pronounced than in the three-dimensional case. For instance, the flux profile of the weak Fe II  $\lambda 6247.562$  line turns out to be in absorption for the 1D+NLTE case.

How large could the impact of UV-overionization and UV-overexcitation on the iron abundance determination in the star considered be? It is obvious that in order to compensate for a weakening of the emergent spectral line profiles in the stellar atmosphere model under consideration, the iron abundance must be increased in comparison with the LTE case. Figure 4

quantifies this conclusion. It shows the LTE versus NLTE iron abundances derived from either Fe I or Fe II lines in the three- and one-dimensional stellar atmospheric models of the metal-poor star HD 140283. As seen in the figure, the NLTE abundance corrections for the Fe I lines are extremely large in the three-dimensional model ( $\sim 0.9$  dex) and a bit smaller in the one-dimensional case ( $\sim 0.6$  dex). On the other hand, it is very important to point out that NLTE effects on the iron abundance are very pronounced for the weak Fe II lines. The four lines used by Nissen et al. (2002) for their iron abundance determinations reveal strikingly large deviations from the LTE. On average, the NLTE abundance corrections for the Fe II lines increase from  $\sim 0.16$  dex in the one-dimensional case to  $\sim 0.4$  dex in the three-dimensional model.

When NLTE effects are taken into account, we find no difference between the mean Fe I-based abundances obtained via the one- and three-dimensional models [ $\log \epsilon(\text{Fe}) \approx 5.77$ ]. Therefore, when NLTE effects are taken into account, the Fe I lines formed in the three-dimensional model are hardly sensitive to granulation inhomogeneities. However, for the Fe II lines the three-dimensional abundance correction (3D+NLTE – 1D+NLTE) is significant—that is, 0.26 dex.




FIG. 3.—Flux profiles of the weak Fe II line at 6247.562 Å (*top panels*) and of the moderately strong Fe II line at 5316.609 Å (*bottom panels*) calculated in the three-dimensional hydrodynamic model of the metal-poor star HD 140283 ( $\log g = 3.7$ ,  $T_{\text{eff}} = 5700$  K, and  $[\text{Fe}/\text{H}] = -2.5$ ). The profiles were synthesized using the iron abundance  $\log \epsilon = 5.0$  corresponding to the metallicity  $[\text{Fe}/\text{H}] = 2.5$ . The left panels show the results for the NLTE case, while the right panels refer to the LTE calculations. Individual thin curves show the computed emergent fluxes for the “granular” and “intergranular” models corresponding to the 50 spatial grid points situated along one of the  $Y$ -directions in the three-dimensional model atmosphere of the metal-poor subgiant HD 140283. The thick solid line with filled circles in each of the panels shows the resulting spatially averaged flux. Fluxes are given in absolute energy units ( $\text{ergs cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$ ). The profiles associated with the dark intergranular lanes of the stellar granulation have a redshift and a lower continuum flux. Bright granules are characterized by a higher continuum flux and a blue line shift. Note that the flux profile of the Fe II line at 6247.562 Å is observed in absorption ( $W > 0$ ), while the synthesized NLTE flux profile turns out to be in emission (calculated  $W < 0$ ).

When the LTE approximation is used, the mean iron abundance obtained via spectrum synthesis of Fe I lines in the three-dimensional model is lower than for the one-dimensional case (cf. Asplund et al. 1999; Nissen et al. 2002). The differences between the one- and three-dimensional determinations depend on the lower excitation potential  $\chi$  of the spectral line under consideration. They range from  $\sim 0.6$  dex at  $\chi \simeq 0$  eV to  $\sim 0.2$  dex at  $\chi \sim 2$  eV, while the mean difference is smaller ( $\sim 0.3$  dex). In contrast, on average the Fe II lines are insensitive to three-dimensional effects when LTE is assumed.

## 8. REVISING THE STELLAR PARAMETERS

There are several discrepancies indicating that the iron abundance obtained in § 7.2 for the star HD 140283 can be considered reliable neither in the one-dimensional case nor in the three-dimensional case. For example, we have (1) a divergence of the average abundances derived from both ionization stages, (2) a correlation of the abundances derived from Fe I lines with the lower level excitation potential, and (3) a large scatter in the abundances derived from Fe II lines (particularly, the weak ones). The elimination of any of such discrepancies cannot be simply achieved in terms of uncertainties in observed equivalent widths, oscillator strengths, inelastic electron collisional rates, and photoionization cross sections. Our numerical experiments show that the typical errors introduced by uncer-

tainties in such quantities do not exceed 0.1 dex (see Shchukina & Trujillo Bueno 2001). The iron abundance error that results from neglecting the UV haze opacity for the metal-poor star HD 140283 is even smaller than 0.1 dex. In our opinion, a reasonable possibility for explaining the reported discrepancies is to revise the stellar parameters of the subgiant HD 140283.

In this section we aim at finding the “best choice” of stellar parameters for the star HD 140283, both for the NLTE and LTE cases. It is important to emphasize from the outset that in our NLTE computations we have neglected inelastic collisions with neutral hydrogen atoms for excitation and ionization. At present, there are no reliable values for the ensuing collisional rates. The often-used classical Drawin (1968) formula leads to very uncertain estimates and to a substantial overestimation of the collisional rates when detailed quantum mechanical and/or laboratory data is available, such as for Li and Na (see Biberman et al. 1987; Lambert 1993; Holweger 1996; Barklem et al. 1998, 2003; Belyayev et al. 1999; Shchukina & Trujillo Bueno 2001; Asplund et al. 2004), which explains why the conclusions of Gratton et al. (1999) have been criticized. Obviously, inelastic collisions with neutral hydrogen atoms would tend to reduce the NLTE effects. Therefore, the difference between the NLTE and LTE iron abundances may be considered as the maximum effect that inelastic collisions might produce. In our opinion, the tentative conclusion of Korn et al. (2003), achieved via a 1D+NLTE




FIG. 4.—Iron abundances derived from Fe I lines (*filled circles*) and from Fe II lines (*stars*) vs. the lower level excitation potential of the chosen spectral lines. The left panels show the results for the three-dimensional model atmosphere of the star HD 140283, while the right panels show results for the one-dimensional model. *Top panels*: NLTE. *Bottom panels*: LTE. The four weak Fe II lines ( $\lambda\lambda 6149.238, 6247.562, 6432.654, \text{ and } 6456.376 \text{ m\AA}$ ) used by Nissen et al. (2002) are marked with squares. The dashed and dotted horizontal lines indicate the mean abundances found from the set of Fe I and Fe II lines, respectively. The corresponding values of the mean abundances and their standard deviation are given inside the panels.

analysis, that hydrogen collisions might be efficient in the atmosphere of HD 140283 needs to be reinvestigated in the light of three-dimensional hydrodynamic models because, as pointed out by these authors, the lower temperatures of the three-dimensional models would alleviate their need of invoking thermalizing collisions to counterbalance photoionization.

Our analysis for the revision of the stellar parameters is based on the following criteria:

1. The average abundances obtained from Fe I and Fe II lines have to be equal.
2. The best-choice solution has to give the minimum mean standard deviation ( $\sigma$ ) for the abundances derived from Fe I and Fe II lines.
3. Abundances have to be independent of the lower excitation potential ( $\chi$ ). We restrict ourselves to revising only the metallicity and the effective temperature. The given surface gravity of the star HD 140283 is reliable, since it has been obtained from accurate *Hipparcos* parallaxes.

Figure 5 helps us to find the best choice of stellar parameters for the star HD 140283. The left panels show the variation of the Fe I and Fe II abundances with metallicity,  $T_{\text{eff}}$ , and  $\log g$  in a suitable grid of one-dimensional model atmospheres. Our results clearly demonstrate that changes in  $T_{\text{eff}}$  seriously affect the Fe I-based abundances. However, such changes in  $T_{\text{eff}}$  are unimportant for abundance determinations from Fe II lines. On the other hand, the Fe II-based abundances are slightly sensitive to the changes in  $\log g$ , which is not the case for the abundances obtained from Fe I lines. In addition, the dependence on metallicity is more pro-

nounced for Fe I lines, while for Fe II lines there is no such variation. It is interesting to note that when LTE is assumed, then only the correlation of the inferred Fe I abundances with  $T_{\text{eff}}$  is present.

### 8.1. Stellar Parameter Determinations: One-dimensional Case

Consider first the 1D+LTE modeling case for the iron lines. Figure 5 (*middle left-hand panel*) and Figure 4 (*bottom right-hand panel*) show that in this case identical abundances from both Fe I and Fe II lines could be obtained around  $T_{\text{eff}}$  close to 5700 K. In fact, our detailed analysis gives  $T_{\text{eff}} \simeq 5670 \text{ K}$  and  $[\text{Fe}/\text{H}] \simeq -2.3$ . With these stellar parameters, Fe II-based abundances turn out to be insignificantly scattered ( $\sigma \sim 0.1 \text{ dex}$ ) around the average value. However, the abundances derived from Fe I lines show a sensitivity to the lower level excitation potential ( $\chi$ ) that is similar to the trend seen in the bottom right-hand panel of Figure 4.

Second, consider the 1D+NLTE modeling case. In order to reach a good agreement between NLTE abundances derived from Fe I and Fe II lines we would have to reduce the effective temperature down to  $T_{\text{eff}} = 5450 \text{ K}$  and increase the stellar metallicity up to  $[\text{Fe}/\text{H}] \simeq -2.1$ . With such stellar parameters the  $\chi$ -dependence of the Fe I abundances vanishes. However, the abundances derived from Fe II lines spread undesirably around the mean value ( $\sigma \sim 0.3 \text{ dex}$ ).

### 8.2. Stellar Parameter Determinations: Three-dimensional Case

In order to achieve a reasonable estimation of the best choice of stellar parameters the curves of Figure 5 (*left panels*) have to




FIG. 5.—Computed abundances vs. the metallicity (*top panels*), vs. the effective temperature (*middle panels*), and vs. the stellar surface gravity (*bottom panels*) for the set of one-dimensional stellar model atmospheres. *Left panels*: Results for iron, with the solid and dashed lines indicating the mean Fe I- and Fe II-based abundances, respectively. *Right panels*: Results for oxygen, with the solid and dashed lines indicating the mean oxygen abundances derived from the infrared triplet at 7772–7775 Å and the oxygen abundance obtained from the forbidden [O I]  $\lambda$ 6300 line. Thick lines in each of the panels correspond to NLTE, while thin lines correspond to LTE. The standard deviations ( $\sigma$ ) for the observed equivalent widths of the O I and [O I] lines are shown in the middle panel. It is also of interest to note that our 1D+LTE calculations show that the lower the effective temperature the larger the discrepancy in the iron abundance derived from Fe I and Fe II lines (see the thin line in the middle left-hand panel).

be corrected for the three-dimensional effects of granulation inhomogeneities. This means that we need to know how the three-dimensional abundance corrections change with  $[\text{Fe}/\text{H}]$ ,  $T_{\text{eff}}$ , and  $\log g$  for both the LTE and NLTE cases. Unfortunately, a suitable grid of three-dimensional models for several effective temperatures and metallicities similar to that provided by Kurucz for the one-dimensional case is not yet available, besides the fact that NLTE computations in three-dimensions are presently very computationally costly. However, we can try to estimate the true stellar parameters of the star HD 140283 via the following strategy, which is inspired by our conclusion that three-dimensional effects are important for the Fe I lines only if LTE is assumed and that three-dimensional effects are relevant for the Fe II lines only when we allow for departures from LTE.

First, let us choose the 3D+LTE modeling approach. As we have seen, if LTE is assumed then the Fe I lines are sensitive to the three-dimensional effects while the Fe II lines are not. Therefore, if the LTE approximation is used it is much safer to revise the stellar parameters via iron abundances derived from Fe II lines. We find  $T_{\text{eff}} \approx 5700$  and  $[\text{Fe}/\text{H}] \approx -2.3$  (see Fig. 4, *right panels*).

Second, consider the 3D+NLTE approach. It is obvious that in this case we better avoid the use of Fe II lines because we have shown that they are sensitive to both three-dimensional

and NLTE effects. However, we have demonstrated also that at least for the available three-dimensional model of the star HD 140283, the NLTE iron abundance derived from Fe I lines is very similar to that obtained via the 1D+NLTE modeling approach. For this reason, we think that we can estimate the true stellar parameters by choosing Fe I lines and a 1D+NLTE modeling approach. We find that the standard deviation  $\sigma$  minimizes at  $T_{\text{eff}} \approx 5600$  K and  $[\text{Fe}/\text{H}] \approx -2$ . Now the mean iron abundance turns out to be  $\log \epsilon(\text{Fe}) = 5.54 \pm 0.14$ . Moreover, the dependence of the derived iron abundances with  $\chi$  becomes negligible.

### 8.3. Uncertainties in the Determination of the Stellar Parameters and the Oxygen Abundance

As we have just seen, depending on the approach used for modeling the iron lines, we get four best choice sets for the stellar parameters of the star HD 140283, with  $T_{\text{eff}}$  in the range  $\approx 5450$ – $5700$  K and  $[\text{Fe}/\text{H}]$  between  $-2.0$  and  $-2.3$  (see Table 4). How large could the impact of such uncertainties in the stellar parameters of the star HD 140283 be on the determination of the oxygen abundance? The right panels of Figure 5 clarify this issue. We can see that when one-dimensional models are used, then both the O I- and [O I]-based abundances are sensitive to changes in  $T_{\text{eff}}$ , but not too much to changes in  $\log g$  and/or in metallicity. The sensitivity of the NLTE abundance corrections

TABLE 4  
STELLAR PARAMETERS OF HD 140283

Case	$T_{\text{eff}}$ (K)	[Fe/H]	$\log g$
1D+LTE .....	5670	-2.3	3.7
1D+NLTE.....	5450	-2.1	3.7
3D+LTE .....	5700	-2.3	3.7
3D+NLTE.....	5600	-2.0	3.7

to these parameters ( $\log g$  and metallicity) is much smaller. On average, they vary between 0.17 and 0.19 dex for O I lines, while it is negligible for the [O I] line ( $<0.01$  dex). Our 1D+NLTE modeling of oxygen lines shows that abundances derived from O I and [O I] lines give the same value [i.e.,  $\log \epsilon(\text{O}) = 7.15$ ] for  $T_{\text{eff}} = 5650$  K. We now point out that among our best choice sets of stellar parameters only one has a  $T_{\text{eff}}$ -value ( $T_{\text{eff}} = 5450$  K), which considerably deviates from that with 5650 K. Such relatively low value for the effective temperature (i.e.,  $T_{\text{eff}} = 5450$  K) causes a large disagreement between the NLTE O I- and [O I]-based abundances amounting to  $\sim 0.34$  dex. Moreover, as we have indicated above, the set of parameters  $T_{\text{eff}} = 5450$  K and  $[\text{Fe}/\text{H}] = -2.1$  produces an unpleasant scatter of the Fe II-based abundances around the mean value. In view of these factors we think reasonable to eliminate such particular set of stellar parameters.

The remaining three sets of stellar parameters, i.e.,  $T_{\text{eff}} = 5670$  K and  $[\text{Fe}/\text{H}] = -2.3$ ,  $T_{\text{eff}} = 5700$  K, and  $[\text{Fe}/\text{H}] = -2.3$ , and  $T_{\text{eff}} = 5600$  K and  $[\text{Fe}/\text{H}] = -2.0$ , correspond to the case of 1D+LTE modeling for both Fe I and Fe II lines, to 3D+LTE modeling of Fe II lines, and to 3D+NLTE modeling of Fe I lines, respectively. It is important to note that this range of effective temperatures is similar to the typical uncertainty in the determination of  $T_{\text{eff}}$  via the application of the infrared flux method. The small differences in the effective temperature implies a rather small oxygen abundance variation. In particular, our 1D+NLTE oxygen abundance determination from the O I IR lines using the former set gives  $\log \epsilon(\text{O}) = 7.13$ , while the latter set leads to  $\log \epsilon(\text{O}) = 7.20$ . For the [O I] line we get  $\log \epsilon(\text{O}) = 7.16$  and  $\log \epsilon(\text{O}) = 7.11$ , respectively.

We must consider now how the dependencies shown for the oxygen lines in the right-hand panels of Figure 5 are going to be modified due to the effect of granulation inhomogeneities. As mentioned above, a major obstacle to investigating this point is the lack of a suitable grid of three-dimensional models. In any case, this three-dimensional issue concerns especially the [O I] line, given its significant sensitivity to granulation inhomogeneities. According to Nissen et al. (2002), the main effect is due to differences in the continuum opacities between the one- and three-dimensional models. Since the opacity is provided primarily by  $\text{H}^-$  atoms the strength of the weak [O I] line turns out to be inversely proportional to electron density. In the three-dimensional hydrodynamic model of the star HD 140283, which has been obtained assuming  $T_{\text{eff}} = 5700$  K and  $[\text{Fe}/\text{H}] = -2.5$ , the deficit of electrons is particularly large in the cool surface layers of the granular regions, while it is less pronounced in both the intergranular plasma and in the one-dimensional model. As a result, the spatially averaged equivalent width of the [O I] line in such a three-dimensional model increases, and consequently the derived oxygen abundance decreases in comparison with the one-dimensional modeling result. Obviously, the three-dimensional correction obtained in § 6.2 for this forbidden line assuming that the stellar parameters are  $T_{\text{eff}} =$

5700 K and  $[\text{Fe}/\text{H}] = -2.5$  cannot be used for the 3D+NLTE case with lower effective temperature ( $T_{\text{eff}} = 5600$  K) and higher metallicity ( $[\text{Fe}/\text{H}] = -2.0$ ).

As pointed out by Asplund et al. (1999), the deficit in the number density of electrons depends mainly on the metallicity. Three-dimensional hydrodynamic simulations for higher  $[\text{Fe}/\text{H}]$  values will have larger surface temperatures and electron densities. Thus, one can expect that at  $[\text{Fe}/\text{H}] = -2$  the three-dimensional abundance correction for the [O I] line has to be smaller. Obviously, the *exact* value of such corrections will have to await the development of a new three-dimensional model for the revised stellar parameters. It should be noted that such a sensitivity to three-dimensional effects may be assumed also for the UV lines of OH. However, the impact on three-dimensional abundance corrections when changing the metallicity from  $-2.5$  to  $-2.0$  seems to be rather small if we suppose that the results obtained by Asplund & García Pérez (2001) for dwarfs can be applied to the subgiant HD 140283 (see their Table 2, which shows that the three-dimensional effects on the UV lines of OH are  $\sim -0.5$  at  $[\text{Fe}/\text{H}] = -2$  and  $\sim -0.6$  at  $[\text{Fe}/\text{H}] = -3$ ). The additional conclusion that follows from their Table 2 is a high sensitivity of the oxygen abundances derived from the OH molecules to the metallicity of the star. As a result, a three-dimensional model with higher metallicity ( $[\text{Fe}/\text{H}] = -2$ ) might bring the oxygen abundances derived from molecules [ $\log \epsilon(\text{O}) = 7.20$  for  $T_{\text{eff}} \approx 5800$ ] to closer agreement with the abundance derived from the O I triplet [ $\log \epsilon(\text{O}) \approx 7.20$  for  $T_{\text{eff}} = 5600$  K.]

An additional problem is related to observations of the [O I] line in the star HD 140283. According to Nissen et al. (2002), this weak line ( $W \simeq 0.5$  mÅ) is blended with a stronger  $\text{H}_2\text{O}$  line ( $W \approx 1$  mÅ). These authors estimated that the available equivalent width measurements have an uncertainty of  $\pm 0.2 - \pm 0.3$  mÅ. Such an uncertainty is very large, because it is about 50% of the observed equivalent width, and implies an uncertainty of  $\pm 0.13 - \pm 0.19$  dex in the derived oxygen abundance (Fig. 5, *middle right-hand panel*).

Alternatively, in a first approximation we could neglect the sensitivity of the O I lines to the granulation inhomogeneities. Furthermore, the observation errors for these lines in HD 140283 are relatively small. For example, the observed equivalent widths of the O I triplet that have been published in the literature (Abia & Rebolo 1989; Tomkin et al. 1992; Nissen et al. 2002; G. Israelian 2003, private communication), and which we have used in our investigation, differ by about  $\pm 1$  mÅ. Their impact on the derived oxygen abundance lies in the range  $\pm 0.06$  dex (Fig. 5, *middle right-hand panel*). It is clear that the O I lines seem to be the most reliable lines for deriving the oxygen abundance in the star HD 140283.

Therefore, when the stellar parameters are revised neglecting three-dimensional and NLTE effects for the iron lines, and when the oxygen abundance derived from the O I lines is corrected for NLTE effects, then the mean value of the oxygen-to-iron ratio seems to be  $[\text{O}/\text{Fe}] \simeq 0.7$  at  $[\text{Fe}/\text{H}] = -2.3$ . This value is in good agreement with the quasi-linear trend of  $[\text{O}/\text{Fe}]$  versus  $[\text{Fe}/\text{H}]$  obtained by Nissen et al. (2002) using standard one-dimensional modeling. However, the key point to keep in mind is that a 3D+NLTE modeling of the O I IR triplet (using the 3D+NLTE stellar parameters derived from the Fe I lines) would give  $[\text{O}/\text{Fe}] \simeq 0.5$  at  $[\text{Fe}/\text{H}] = -2.0$ . Obviously, the analysis of a single metal-poor star is not enough to opt for one of the following two possibilities: (1) a linear rise in the  $[\text{O}/\text{Fe}]$  ratio versus metallicity (see, e.g., Fig. 11 of Israelian et al. 2001) or (2) a plateau between 0.4 and 0.5 dex in the metallicity range

$-3 < [\text{Fe}/\text{H}] < -1$ .<sup>4</sup> However, we think that our investigation indicates the “road to be taken” toward the resolution of the problem.

## 9. CONCLUSIONS

We have derived the solar oxygen abundance from the O I IR triplet using the Asplund et al. (2000) three-dimensional hydrodynamic model of the solar photosphere and a NLTE modeling approach. We find  $\log \epsilon(\text{O}) = 8.70 \pm 0.06$ , which is in good agreement with the determinations of Allende Prieto et al. (2001), Shchukina et al. (2003), and Asplund et al. (2004). Actually, if for the O I IR triplet we use the same oscillator strengths adopted by Asplund et al. (2004), we then obtain a value very similar to that reported by them [i.e.,  $\log \epsilon(\text{O}) = 8.65$ ].

In agreement with previous investigations, we confirm that LTE synthesis of Fe I lines in the three-dimensional hydrodynamic model of this star gives flux profiles significantly different from those obtained in a one-dimensional model for the same stellar parameters. However, when the LTE approximation is used for the synthesis of Fe II lines we find negligible differences between the one- and three-dimensional cases.

The differences between the iron abundances obtained from Fe I lines when assuming LTE in the one-dimensional and three-dimensional models of the star HD 140283 turn out to depend on the lower excitation potential of the spectral line under consideration. The NLTE abundance corrections for Fe I lines are significant, especially for the low-excitation lines in the three-dimensional model.

We have three particularly important results concerning the determination of the iron abundance in metal-poor stars like HD 140283:

1. If NLTE effects are taken into account when synthesizing the Fe I lines in the one- and three-dimensional models of this star, then the derived iron abundance turns out to be practically the same in both cases.
2. Contrary to a generally accepted belief, the Fe II lines turn out to be significantly affected by NLTE effects. In particular, a full NLTE modeling should be carried out for the Fe II lines with

upper levels of odd parity starting from  $z^6D^o$ , and especially for the weaker lines.

3. The iron abundance of the star HD 140283 is estimated to be  $\sim 0.5$  dex higher than previously thought.

For the O I IR triplet the NLTE correction of the oxygen abundance in the one- and three-dimensional models is practically the same and less than 0.2 dex. Such a NLTE correction is negligible for the [O I] line. We have confirmed that the mean value of the oxygen abundance derived from the O I IR lines is insensitive to three-dimensional effects, while such effects reduce the abundance derived from the [O I] line by  $\sim 0.2$  dex.

With the NLTE calculations presented here for Fe and O, both the one- and three-dimensional models lead to inconsistencies in the iron and oxygen abundance determinations. The discrepancies in the derived iron and oxygen abundances cannot be removed by taking into account NLTE and/or three-dimensional effects. This result has led us to investigate whether these discrepancies can be resolved by a modification of the stellar parameters of the star HD 140283. We have shown that previous studies may have underestimated the metallicity of this star and overestimated its effective temperature. We find  $[\text{Fe}/\text{H}] \simeq -2.0$  (instead of  $[\text{Fe}/\text{H}] \simeq -2.5$ ) and  $T_{\text{eff}} = 5600$  (instead of  $T_{\text{eff}} = 5700$ ). With these new stellar parameters the iron and oxygen abundances in the star HD 140283 would be  $\log \epsilon(\text{Fe}) = 5.54 \pm 0.14$  and  $\log \epsilon(\text{O}) = 7.20 \pm 0.06$ , respectively. Taking into account the low value for the solar oxygen abundance [i.e.,  $\log \epsilon_{\odot}(\text{O}) = 8.70 \pm 0.06$ ] and our previously determined value of the solar iron abundance [i.e.,  $\log \epsilon_{\odot}(\text{Fe}) = 7.50 \pm 0.10$ ; see Shchukina & Trujillo Bueno 2001], we find that the oxygen-to-iron abundance ratio is  $[\text{O}/\text{Fe}] \approx 0.5$  at  $[\text{Fe}/\text{H}] = -2$ .

Besides the reported NLTE effects for the Fe II lines, our main conclusion here is that the metallicity of this type of metal-poor stars might well be significantly larger than previously thought. Obviously, the analysis of a single metal-poor star is not enough to fully resolve the puzzling behavior of the O/Fe ratio in metal-poor stars. However, we think that the present investigation at least indicates the road to be taken.

We are very grateful to Ramón García López and Garik Israelian for allowing us to use their stellar observations of iron and oxygen lines. We are also grateful to I. Vasiljeva for helping us with the radiative transfer computations during the early stages of this project. This work has been funded by the European Commission through INTAS grant 00-00084 and by the Spanish Ministerio de Educación y Ciencia through project AYA2004-05792.

<sup>4</sup> We point out that with the stellar parameters used by Asplund et al. (1999) for the three-dimensional modeling of the star HD 140283 (i.e.,  $T_{\text{eff}} = 5700$  and  $[\text{Fe}/\text{H}] = -2.5$ ), the NLTE oxygen abundance [ $\log \epsilon(\text{O}) = 7.08 \pm 0.06$ ] derived from the O I IR triplet and the NLTE Fe I-based iron abundance [ $\log \epsilon(\text{Fe}) = 5.77 \pm 0.186$ ] leads to a small oxygen-to-iron ratio ( $[\text{O}/\text{Fe}] \sim 0.1$  dex). Moreover, there is no overabundance of oxygen at all if one uses the [O I]-based abundance. It is also of interest to mention that in our preliminary study (see Shchukina et al. 2003) we obtained a slightly larger value for  $[\text{O}/\text{Fe}]$  because we used a smaller number of Fe I lines.

## REFERENCES

- Abia, C., & Rebolo, R. 1989, *ApJ*, 347, 186  
 Allende Prieto, C., García López, R. J., Lambert, D. L., & Gustafsson, B. 1999, *ApJ*, 526, 991  
 Allende Prieto, C., Lambert, D. L., & Asplund, M. 2001, *ApJ*, 556, L63  
 Alcock, R. C. 1968, *Sol. Phys.*, 5, 260  
 Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197  
 Anstee, S. D., & O’Mara, B. J. 1995, *MNRAS*, 276, 859  
 Asplund, M., & García Pérez, A. E. 2001, *A&A*, 372, 601  
 Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., & Kiselman, D. 2004, *A&A*, 417, 751  
 Asplund, M., Ludwig, H.-G., Nordlund, Å., & Stein, R. F. 2000, *A&A*, 359, 669  
 Asplund, M., Nordlund, Å., Trampedach, R., & Stein, R. F. 1999, *A&A*, 346, L17  
 Athay, R. G., & Lites, B. W. 1972, *ApJ*, 176, 809  
 Barbuy, B. 1988, *A&A*, 191, 121  
 Barklem, P. S., Belyaev, A.K., & Asplund, M. 2003, *A&A*, 409, L1  
 Barklem, P. S., & O’Mara, B. J. 1997, *MNRAS*, 290, 102  
 Barklem, P. S., O’Mara, B. J., & Ross, J. E. 1998, *MNRAS*, 296, 1057  
 Bautista, M. A. 1997, *A&AS*, 122, 167  
 Belyaev, A., Grosser, J., Hahne, J., & Menzel, T. 1999, *Phys. Rev. A*, 60, 2151  
 Biberian, L. M., Vorobjev, V. S., & Yakubov, I. T. 1987, *Kinetics of Non-equilibrium Low-Temperature Plasma* (New York: Plenum)  
 Boesgaard, A. M., King, J. R., Deliyannis, C. P., & Vogt, S. 1999, *AJ*, 117, 492  
 Carlsson, M., & Judge, P. 1993, *ApJ*, 402, 344  
 Carretta, E., Gratton, R. G., & Sneden, C. 2000, *A&A*, 356, 238  
 Cavallo, R. M., Pilachowski, C. A., & Rebolo, R. 1997, *PASP*, 109, 226  
 Cram, L. E., Rutten, R. J., & Lites, B. W. 1980, *ApJ*, 241, 374  
 Delbouille, L., Neven, L., & Roland, G. 1973, *Photometric Atlas of the Solar Spectrum from  $\lambda 3000$  to  $\lambda 10,000$*  (Liege: Inst. d’Astrophys, Univ. de Liege)

- Drawin, H. W. 1968, *Z. Phys.*, 211, 404
- Fuhr, J. R., Martin, G. A., & Wiese, W. L. 1988, *J. Phys. Chem. Ref. Data*, 17, 24
- Fulbright, J., & Johnson, J. 2003, *ApJ*, 595, 1154
- Fulbright, J., & Kraft, R. 1999, *AJ*, 118, 527
- Gratton, R. G., Caretta, E., Eriksson, K., & Gustafsson, B. 1999, *A&A*, 350, 955
- Grevesse, N., & Sauval, A. J. 1998, *Space Sci. Rev.*, 85, 161
- Grevesse, N., Sauval, A. J., & van Dishoeck, E. F. 1984, *A&A*, 141, 10
- Gurtovenko, E. A., & Kostik, R. I. 1989, *Fraunhofer Spectrum and the System of Solar Oscillator Strengths* (Russian language ed.; Kiev: Naukova Dumka)
- Gustafsson, B., Bell, R. A., Eriksson, K., & Nordlund, Å. 1975, *A&A*, 42, 407
- Holweger, H. 1996, *Phys. Scr.*, T65, 151
- . 2001, in *AIP Conf. Proc.* 598, *Solar and Galactic Composition*, ed. R. F. Wimmer-Schweingruber (Melville: AIP), 23
- Holweger, H., & Müller, E. 1974, *Sol. Phys.*, 39, 19
- Israelian, G., García López, R. J., & Rebolo, R. 1998, *ApJ*, 507, 805
- Israelian, G., Rebolo, R., García López, R. J., Bonifacio, P., Molaro, P., Basri, G., & Shchukina, N. 2001, *ApJ*, 551, 833
- Israelian, G., Shchukina, N., Rebolo, R., Basri, G., González Hernández, J. I., & Kajino, T. 2004, *A&A*, 419, 1095
- King, J. R., & Boesgaard, A. M. 1995, *AJ*, 109, 383
- Kiselman, D. 1993, *A&A*, 275, 269
- . 2001, *NewA Rev.*, 45, 559
- Kiselman, D., & Nordlund, Å. 1995, *A&A*, 302, 578
- Korn, A. J., Shi, J., & Gehren, T. 2003, *A&A*, 407, 691
- Kostik, R. I., Shchukina, N. G., & Rutten, R. J. 1996, *A&A*, 305, 325
- Kraft, R. P., Sneden, C., Langer, G. E., & Prosser, C. F. 1992, *AJ*, 104, 645
- Kurucz, R. L. 1993, *Kurucz CD-Rom 13, ATLAS9 Stellar Atmosphere Programs and 2km/s Grid* (Cambridge: SAO)
- Kurucz, R. L., & Bell, B. 1995, *Kurucz CD-ROM 23, Atomic Line List* (Cambridge: SAO)
- Lambert, D. L. 1978, *MNRAS*, 182, 249
- . 1993, *Phys. Scr.*, T47, 186
- Meléndez, J., & Barbuy, B. 2002, *ApJ*, 575, 474
- Mishenina, T., Korotin, S., Klochkova, V., & Panchuk, V. 2000, *A&A*, 353, 978
- Moore, Ch. 1959, *A Multiplet Table of Astrophysical Interest* (NBS Tech. Note; Washington, DC: NBS)
- Müller, E., Baschek, B., & Holweger, H. 1968, *Sol. Phys.*, 3, 125
- Nissen, P. E., Primas, F., Asplund, M., & Lambert, D. L. 2002, *A&A*, 390, 235
- Rutten, R. J. 1988, in *IAU Colloq. 94, Physics of Formation of Fe II Lines Outside LTE*, ed. R. Viotti, A. Vittone, & M. Friedjung (Dordrecht: Reidel), 185
- Sauval, A. J., Grevesse, N., Brault, J. B., Stokes, G. M., & Zander, R. 1984, *ApJ*, 282, 330
- Shchukina, N. G. 1987, *Kinemat. Phys. Celest. Bodies*, 3, 36
- Shchukina, N., & Trujillo Bueno, J. 2001, *ApJ*, 550, 970
- Shchukina, N. G., Vasiljeva, I. E., Trujillo Bueno, J., & Asplund, M. 2003, in *IAU Symp. 210, Modeling of Stellar Atmospheres*, ed. N. Piskunov, W. W. Weiss, & D. F. Gray (San Francisco: ASP), 10
- Socas-Navarro, H., & Trujillo Bueno, J. 1997, *ApJ*, 490, 383
- Stein, R. F., & Nordlund, Å. 1998, *ApJ*, 499, 914
- Takeda, Y. 2003, *A&A*, 402, 343
- Thévenin, F., & Idiart, T. 1999, *ApJ*, 521, 753
- Tomkin, J., Lemke, M., Lambert, D. L., & Sneden, C. 1992, *AJ*, 104, 1568
- Trujillo Bueno, J. 2003, in *ASP Conf. Ser. 288, Stellar Atmosphere Modeling*, ed. I. Hubeny, D. Mihalas, & K. Werner (San Francisco: ASP), 551
- Trujillo Bueno, J., & Fabiani Bendicho, P. 1995, *ApJ*, 455, 646